

11th International Conference on Hard and Electromagnetic Probes of High-Energy Nuclear Collisions



Beitrag ID: 300

Typ: Talk

Spin-Induced Interactions and Heavy-Quark Transport in the QGP

Dienstag, 28. März 2023 14:40 (20 Minuten)

We extend a previously constructed T-matrix approach for heavy quarks in the quark-gluon plasma (QGP) to include inverse-mass ($1/M$) corrections, i.e. the spin-orbit, spin-spin and tensor forces, between partons. Based on the vacuum Cornell potential as the interaction kernel for the T-matrix equation, we first confirm that the experimental charmonium and bottomonium spectroscopy in vacuum are much improved by employing a confining potential that is a mixture of vector and scalar interactions, rather than a purely scalar one. We then apply the refined potential to calculate the in-medium single-parton spectral functions at finite temperature self-consistently and constrained by various thermal lattice-QCD data. Finally, we study the consequences for the in-medium charm-quark transport coefficients at different temperatures. It turns out that the mixing effect for confining potential significantly enhances the friction coefficient, $A(p)$, for charm quarks in the QGP over previous calculations with a purely scalar potential. Our results may have significant implications for the microscopic description of heavy-flavor transport in heavy-ion collisions at RHIC and the LHC.

Experiment/Theory

Theory/Phenomenology

Affiliation

Cyclotron Institute and Department of Physics and Astronomy, Texas A&M University

Hauptautor: TANG, Zhanduo

Co-Autor: Dr. RAPP, Ralf (Cyclotron Institute and Department of Physics and Astronomy, Texas A&M University)

Vortragende(r): TANG, Zhanduo

Sitzung Einordnung: Parallel: Heavy Flavours & Quarkonia

Track Klassifizierung: Heavy flavor and quarkonia