

11th International Conference on Hard and Electromagnetic Probes of High-Energy Nuclear Collisions



Beitrag ID: 81

Typ: Talk

Electroweak-boson measurements from small to large collision systems with ALICE at the LHC

Dienstag, 28. März 2023 17:30 (20 Minuten)

Electroweak bosons produced in hard-scattering processes at the early stage of the collision, are efficient probes of the initial state of the collision. While the W measurements in pp collisions are a stringent test of perturbative QCD-based calculations and production mechanisms, they can constrain the nuclear parton distribution functions in p-Pb and Pb-Pb collisions.

Electroweak bosons are studied with ALICE in pp collisions at $\sqrt{s} = 13$ TeV, p-Pb collisions at $\sqrt{s_{NN}} = 8.16$ TeV and Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV via their leptonic decays in the muon and electron channels at forward rapidity ($-4.0 < \eta < -2.5$) and midrapidity ($|\eta| < 0.8$), respectively. The observations in p-Pb and Pb-Pb collisions at forward rapidity give access to low Bjorken- x values, a phase-space region poorly constrained by heavy-ion experiments.

The latest W-boson results concerning differential measurements of the normalised production yields, production cross sections, nuclear modification factors and lepton-charge asymmetry as a function of rapidity, transverse momentum, collision centrality and charged-particle multiplicity are presented. The production of W bosons in association with hadrons as a function of the charged-particle multiplicity in pp collisions is reported as well. Comparisons with model calculations are discussed.

Experiment/Theory

ALICE

Affiliation

ALICE

Hauptautor: SAKAI, Shingo (Univ. of Tsukuba)

Vortragende(r): SAKAI, Shingo (Univ. of Tsukuba)

Sitzung Einordnung: Parallel: Electromagnetic & Electroweak Probes

Track Klassifizierung: Electromagnetic and electroweak probes