



Beitrag ID: 367

Typ: Talk

## Disentangle effects from the initial stage and the evolution stage in heavy ion collisions using EPOS and PHSD

*Freitag, 31. März 2023 09:00 (5 Minuten)*

Ultrarelativistic heavy-ion collisions at RHIC and the LHC provide a hot and ultra-dense form of matter composed of deconfined quarks and gluons, named QGP. Different models like EPOS and PHSD allow to study the space-time evolution of such heavy-ion collisions. Their dynamics is complicated; hence, various stages should be considered. The first is the primary scattering which defines to a large extent the matter distribution in the phase-space. The second stage concerns the evolution of the partonic system until the system is sufficiently dilute to hadronize. The EPOSi+PHSD approach is introduced in this thesis, in which the EPOS model is used to determine the initial distribution of matter (partons/hadrons). This part is referred to as EPOSi. Then PHSD is employed to simulate the evolution of the matter in a non-equilibrium transport approach, referred to as PHSD<sub>e</sub>. The coupling of the two approaches is non-trivial and not straight-forward. Comparing the three models, EPOS, EPOSi+PHSD<sub>e</sub>, and PHSD, interesting results find concerning their respective space-time evolutions. The results demonstrate considerably different behavior in terms of radial expansion, especially asymmetric expansion, indicating that these three models will provide different results concerning key observables (  $p_T/m_T$  spectra,  $y/\eta$  distribution,  $v_2/3/4$ ) for Au-Au collisions at 200 GeV/A.

### Experiment/Theory

### Affiliation

**Hauptautor:** JAFARPOUR, Mahbobeh

**Vortragende(r):** JAFARPOUR, Mahbobeh

**Sitzung Einordnung:** Plenary Session VIII